**Recognition, Causes and Interventions** 

August 27, 2014

Stephen M. Soreff, MD

**Education Initiatives, LLC** 

e-mail <u>soreffs15@aol.com</u> <u>http://eduinitiatives.net</u>

**Telephone 603 895-6120** 




**Recognition, Causes and Interventions** 




- Statistics
- Definitions
- Recognition –Signs and Symptoms
- Causes
- Interventions
- Post-Interventions

**Recognition, Causes and Interventions** 



# Adolescent Suicide: Recognition, Causes and Interventions



**Recognition, Causes and Interventions** 

### Suicide killing oneself

A completed suicide

A suicide attempt

A suicide gesture

A suicide gamble

A suicide equivalent

**Recognition, Causes and Interventions** 

#### Recognition –Signs and Symptoms










**Recognition, Causes and Interventions** 

- Suicidal individuals have a number of characteristics, including the following:
- A preoccupation with death
- A sense of isolation and withdrawal
- Few friends or family members
- An emotional distance from others
- Distraction and lack of humor They often seem to be "in their own world" and lack a sense of humor (anhedonia)
- Focus on the past They dwell on past losses and defeats and anticipate no future; they voice the notion that others and the world would be better off without them
- Haunted and dominated by hopelessness and helplessness

# Adolescent Suicide: Recognition, Causes and Interventions



http://blogs.cornell.edu/into2040/2011/09/21/the-clustering-coefficient/ 7/27/2014

#### H<sub>x</sub>2 =Hopelessness and Helplessness

I can't pretend that I'm okay anymore, I can't fake it anymore, I'm hopeless, helpless and im afraid I'll never get better.

**Recognition, Causes and Interventions** 

### Numerous activities are associated with suicidal potential, including the following:


- Making a will
- Getting the house and affairs together
- Unexpectedly visiting friends and family members
- Purchasing a gun, hose, or rope
- Writing a suicide note
- Visiting a primary care physician A significant number of people see their primary care physician within 3 weeks before they commit suicide

#### **Recognition, Causes and Interventions**

#### **Correlations, Relationships and Causes**

- Sex
- Drugs and Alcohol
- Culture/Nation/Religion
- Location Rural/Urban
- Mental Illness/Depression
- Loss of Perspectives
- Bullying
- Media/Contagion
- Family History
- Season
- Anomie
- Medical/Biologic
- Ambivalence
- Criminal-justice system
- Doctor visit


Recognition, Causes and Interventions



SOURCE: Centers for Disease Control and Prevention

### Adolescent Suicide: Loss of Perspective





# Adolescent Suicide: Bullying





#### Ambivalence (n):

The state of having simultaneous, sometimes conflicting feelings towards something – like feeling happy and sad at the same time.






**Recognition, Causes and Interventions** 

#### **Interventions**

- A. Immediacy
- B. Be with the adolescent at all times-do not leave the person alone
- C. Multiple simultaneous methods the danger of assumptions
- D. Search of the person
- E. Definite, defined follow-up
- F. Involvement of significant others
- G. Homicide

**Recognition, Causes and Interventions** 

## Who young people turn to for help these days





Recognition, Causes and Interventions

#### **Post-vention**

- A. Meet with family
- B. Listen
- C. Express your concerns

**Recognition, Causes and Interventions** 



Suicide represents a permanent solution to a temporary problem.

Suicide can be prevented.

#### YOU CAN MAKE THE DFFERENCE



Ask

Listen

with your third ear and gut

Act